

DSM-5 Diagnoses and New ICD-10-CM Codes

As Ordered in the DSM-5 Classification



| Disorder | DSM-5 Recommended ICD-10-CM Code for use through September 30, 2017 | DSM-5 Recommended ICD-10-CM Code for use beginning October 1, 2017 |
|--|---|--|
| Avoidant/Restrictive Food Intake Disorder | F50.89 | F50.82 |
| Alcohol Use Disorder, Mild | F10.10 | F10.10 |
| Alcohol Use Disorder, Mild, in early or sustained remission | F10.10 | F10.11 |
| Alcohol Use Disorder, Moderate | F10.20 | F10.20 |
| Alcohol Use Disorder, Moderate, in early or sustained remission | F10.20 | F10.21 |
| Alcohol Use Disorder, Severe | F10.20 | F10.20 |
| Alcohol Use Disorder, Severe, in early or sustained remission | F10.20 | F10.21 |
| Cannabis Use Disorder, Mild | F12.10 | F12.10 |
| Cannabis Use Disorder, Mild, in early or sustained remission | F12.10 | F12.11 |
| Cannabis Use Disorder, Moderate | F12.20 | F12.20 |
| Cannabis Use Disorder, Moderate, in early or sustained remission | F12.20 | F12.21 |
| Cannabis Use Disorder, Severe | F12.20 | F12.20 |
| Cannabis Use Disorder, Severe, in early or sustained remission | F12.20 | F12.21 |
| Phencyclidine Use Disorder, Mild | F16.10 | F16.10 |
| Phencyclidine Use Disorder, Mild, in early or sustained remission | F16.10 | F16.11 |
| Phencyclidine Use Disorder, Moderate | F16.20 | F16.20 |
| Phencyclidine Use Disorder, Moderate, in early or sustained remission | F16.20 | F16.21 |
| Phencyclidine Use Disorder, Severe | F16.20 | F16.20 |
| Phencyclidine Use Disorder, Severe, in early or sustained remission | F16.20 | F16.21 |
| Other Hallucinogen Use Disorder, Mild | F16.10 | F16.10 |
| Other Hallucinogen Use Disorder, Mild, in early or sustained remission | F16.10 | F16.11 |
| Other Hallucinogen Use Disorder, Moderate | F16.20 | F16.20 |
| Other Hallucinogen Use Disorder, Moderate, in early or sustained remission | F16.20 | F16.21 |
| Other Hallucinogen Use Disorder, Severe | F16.20 | F16.20 |
| Other Hallucinogen Use Disorder, Severe, in early or sustained remission | F16.20 | F16.21 |
| Inhalant Use Disorder, Mild | F18.10 | F18.10 |
| Inhalant Use Disorder, Mild, in early or sustained remission | F18.10 | F18.11 |
| Inhalant Use Disorder, Moderate | F18.20 | F18.20 |

| | | |
|---|---------|---------|
| Inhalant Use Disorder, Moderate, in early or sustained remission | F18.20 | F18.21 |
| Inhalant Use Disorder, Severe | F18.20 | F18.20 |
| Inhalant Use Disorder, Severe, in early or sustained remission | F18.20 | F18.21 |
| Opioid Use Disorder, Mild | F11.10 | F11.10 |
| Opioid Use Disorder, Mild, in early or sustained remission | F11.10 | F11.11 |
| Opioid Use Disorder, Moderate | F11.20 | F11.20 |
| Opioid Use Disorder, Moderate, in early or sustained remission | F11.20 | F11.21 |
| Opioid Use Disorder, Severe | F11.20 | F11.20 |
| Opioid Use Disorder, Severe, in early or sustained remission | F11.20 | F11.21 |
| Sedative, Hypnotic, or Anxiolytic Use Disorder, Mild | F13.10 | F13.10 |
| Sedative, Hypnotic, or Anxiolytic Use Disorder, Mild, in early or sustained remission | F13.10 | F13.11 |
| Sedative, Hypnotic, or Anxiolytic Use Disorder, Moderate | F13.20 | F13.20 |
| Sedative, Hypnotic, or Anxiolytic Use Disorder, Moderate, in early or sustained remission | F13.20 | F13.21 |
| Sedative, Hypnotic, or Anxiolytic Use Disorder, Severe | F13.20 | F13.20 |
| Sedative, Hypnotic, or Anxiolytic Use Disorder, Severe, in early or sustained remission | F13.20 | F13.21 |
| Amphetamine-type Substance Use Disorder, Mild | F15.10 | F15.10 |
| Amphetamine-type Substance Use Disorder, Mild, in early or sustained remission | F15.10 | F15.11 |
| Amphetamine-type Substance Use Disorder, Moderate | F15.20 | F15.20 |
| Amphetamine-type Substance Use Disorder, Moderate, in early or sustained remission | F15.20 | F15.21 |
| Amphetamine-type Substance Use Disorder, Severe | F15.20 | F15.20 |
| Amphetamine-type Substance Use Disorder, Severe, in early or sustained remission | F15.20 | F15.21 |
| Cocaine Use Disorder, Mild | F14.10 | F14.10 |
| Cocaine Use Disorder, Mild, in early or sustained remission | F14.10 | F14.11 |
| Cocaine Use Disorder, Moderate | F14.20 | F14.20 |
| Cocaine Use Disorder, Moderate, in early or sustained remission | F14.20 | F14.21 |
| Cocaine Use Disorder, Severe | F14.20 | F14.20 |
| Cocaine Use Disorder, Severe, in early or sustained remission | F14.20 | F14.21 |
| Tobacco Use Disorder, Moderate | F17.200 | F17.200 |
| Tobacco Use Disorder, Moderate, in early or sustained remission | F17.200 | F17.201 |
| Tobacco Use Disorder, Severe | F17.200 | F17.200 |

| | | |
|--|---------|---------|
| Tobacco Use Disorder, Severe, in early or sustained remission | F17.200 | F17.201 |
| Other (or Unknown) Substance Use Disorder, Mild | F19.10 | F19.10 |
| Other (or Unknown) Substance Use Disorder, Mild, in early or sustained remission | F19.10 | F19.11 |
| Other (or Unknown) Substance Use Disorder, Moderate | F19.20 | F19.20 |
| Other (or Unknown) Substance Use Disorder, Moderate, in early or sustained remission | F19.20 | F19.21 |
| Other (or Unknown) Substance Use Disorder, Severe | F19.20 | F19.20 |
| Other (or Unknown) Substance Use Disorder, Severe, in early or sustained remission | F19.20 | F19.21 |